

ASSOCIATED PRIMES OF LOCAL COHOMOLOGY MODULES OF WEAKLY LASKERIAN MODULES

KAMRAN DIVAANI-AAZAR AND AMIR MAFI

ABSTRACT. The notion of weakly Laskerian modules was introduced recently by the authors. Let R be a commutative Noetherian ring with identity, $\mathfrak{a}$ an ideal of R , and M a weakly Laskerian module. It is shown that if $\mathfrak{a}$ is principal, then the set of associated primes of the local cohomology module $H_{\mathfrak{a}}^i(M)$ is finite for all $i \geq 0$. We also prove that when R is local, then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite for all $i \geq 0$ in the following cases: (1) $\dim R \leq 3$, (2) $\dim R/\mathfrak{a} \leq 1$, (3) M is Cohen-Macaulay and for any ideal $\mathfrak{b}$, with $l = \text{grade}(\mathfrak{b}, M)$, $\text{Hom}_R(R/\mathfrak{b}, H_{\mathfrak{b}}^{l+1}(M))$ is weakly Laskerian.

1. INTRODUCTION

Throughout, let R be a commutative Noetherian ring with identity. Let $\mathfrak{a}$ be an ideal of R and M an R -module. For each $i \geq 0$, the i -th local cohomology module of M with respect to $\mathfrak{a}$ is defined as

$$H_{\mathfrak{a}}^i(M) = \varinjlim_n \text{Ext}_R^i(R/\mathfrak{a}^n, M).$$

The reader can refer to [3] for basic properties of local cohomology. In [7], Hartshorne defined an R -module M to be $\mathfrak{a}$ -cofinite if $\text{Supp}_R M \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(R/\mathfrak{a}, M)$ is finitely generated for all $i \geq 0$. He then asked when the local cohomology modules of a finitely generated module are $\mathfrak{a}$ -cofinite. In this regard, the best known result is that for a finitely generated R -module M if either $\mathfrak{a}$ is principal or R is local and $\dim R/\mathfrak{a} = 1$, then the $H_{\mathfrak{a}}^i(M)$'s are $\mathfrak{a}$ -cofinite. These results are proved in [9, Theorem 1] and [4, Theorem 1], respectively.

It is easy to see that an $\mathfrak{a}$ -cofinite module has only finitely many associated primes. Huneke [8] raised the following question: If M is a finitely generated R -module, then the set of associated primes of $H_{\mathfrak{a}}^i(M)$ is finite for all ideals $\mathfrak{a}$ of R and all $i \geq 0$.

2000 *Mathematics Subject Classification.* 13D45, 13E99.

Key words and phrases. Local cohomology, associated prime ideals, cofiniteness, weakly Laskerian modules, spectral sequences.

Singh [16] gives a counter-example to this question (also see [17] for some other counter-examples). On the other hand, Brodmann and Lashgari [2, Theorem 2.2] showed that if for a finitely generated R -module M and an integer t , the local cohomology module $H_{\mathfrak{a}}^i(M)$ is finitely generated for all $i < t$, then $\text{Ass}_R(H_{\mathfrak{a}}^t(M))$ is finite. For another proof of this result, see [10]. As in [6], an R -module M is called weakly Laskerian if $\text{Ass}_R(M/N)$ is finite for all submodules N . Let M be a weakly Laskerian module and $t \in \mathbb{N}$. In [6], we proved the set of associated primes of the first non $\mathfrak{a}$ -cofinite local cohomology module of M is finite. That clearly implies the result mentioned above due to Brodmann and Lashgari. Also regarding the Artinianness of local cohomology, it is shown in [5], that in many cases weakly Laskerian modules behave similar to finitely generated modules.

In this article, we continue studying the set of associated primes of local cohomology of weakly Laskerian modules. Example 2.3 shows that the class of weakly Laskerian modules is much larger than that of Noetherian modules. Our main aim in this paper is to show that in conjunction with finiteness properties of local cohomology, in many cases weakly Laskerian modules behave similar to finitely generated modules. By applying the known techniques we can get the analogues of some finiteness results for finitely generated modules for weakly Laskerian modules.

In section 2, we first review weakly Laskerian modules. Then we introduce the notion of weak cofiniteness and show that many results concerning cofiniteness of finitely generated modules are valid for weak cofiniteness of weakly Laskerian modules. In particular, it is proved that the Change of Ring Principle for weak cofiniteness holds.

The main results of this article appear in section 3. We prove that the local cohomology modules of a weakly Laskerian module are weakly cofinite in several cases. As a result, we deduce that the sets of associated primes of the local cohomology modules of a weakly Laskerian module are finite in these cases. For instance, it is shown that if $\mathfrak{a}$ is a principal ideal of R and M a weakly Laskerian R -module, then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite for all $i \geq 0$. Also, we prove that if R is a local ring and M a weakly Laskerian R -module, then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite for all $i \geq 0$, in the following cases:

- a) $\dim R \leq 3$,
- b) $\dim R/\mathfrak{a} \leq 1$.

Finally, we show that if M is a Cohen-Macaulay module over a local ring R such

that for any ideal $\mathfrak{b}$ with $l = \text{grade}(\mathfrak{b}, M)$, $\text{Hom}_R(R/\mathfrak{b}, H_{\mathfrak{b}}^{l+1}(M))$ is weakly Laskerian, then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite for all $i \geq 0$. This extends the main result of [1].

2. WEAKLY COFINITE MODULES

In this section, we first recall the definition of weakly Laskerian modules. Then we bring a lemma which is needed in the sequel.

Definition 2.1. i) An R -module M is said to be *Laskerian* if any submodule of M is an intersection of a finite number of primary submodules.
 ii) (See [6]) An R -module M is said to be *weakly Laskerian* if the set of associated primes of any quotient module of M is finite.

Obviously, any Noetherian module is Laskerian and it is clear that any Laskerian module is weakly Laskerian. So, any Noetherian module is weakly Laskerian. We need to the following lemma in the sequel.

Lemma 2.2. i) Let $0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$ be an exact sequence of R -modules. Then M is weakly Laskerian if and only if L and N are both weakly Laskerian. Thus any subquotient of a weakly Laskerian module as well as any finite direct sum of weakly Laskerian modules is weakly Laskerian.
 ii) Let M and N be two R -modules. If M is weakly Laskerian and N is finitely generated, then $\text{Ext}_R^i(N, M)$ and $\text{Tor}_i^R(N, M)$ are weakly Laskerian for all $i \geq 0$.
 iii) Let M be an R -module such that $\text{Supp}_R M$ is finite. Then M is weakly Laskerian. In particular, any Artinian R -module is weakly Laskerian.

Proof. The proof of i) is easy and we leave it to the reader.

ii) See [6, Lemma 2.3].

iii) Let N be a submodule of M . We have

$$\text{Ass}_R(M/N) \subseteq \text{Supp}_R(M/N) \subseteq \text{Supp}_R M.$$

Thus the set of associated primes of any quotient module of M is finite. $\square$

Now, we provide several examples for this class of modules.

Example 2.3. i) An R -module M is said to be *minimax* if M has a finitely generated submodule S such that M/S is Artinian (see [18]). By Lemma 2.2, it follows that any minimax R -module is weakly Laskerian.

ii) Let E be the minimal injective cogenerator of R . If for an R -module M the natural map from M to $\text{Hom}_R(\text{Hom}_R(M, E), E)$ is an isomorphism, then M is said to be Matlis reflexive. Also, an R -module M is said to be linearly compact if each system of congruences

$$x \equiv x_i(M_i),$$

indexed by a set I and where the M_i are submodules of M , has a solution x whenever it has a solution for every finite subsystem. It is known that if either M is reflexive or linearly compact, then M has a finitely generated submodule S such that M/S is Artinian (see e.g. [6, Example 2.2]). Hence any reflexive R -module and also any linearly compact R -module is weakly Laskerian.

Next, we present a generalization of the notion of cofiniteness.

Definition 2.4. Let $\mathfrak{a}$ be an ideal of R and M an R -module. We say that M is $\mathfrak{a}$ -weakly cofinite if $\text{Supp}_R M \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(R/\mathfrak{a}, M)$ is weakly Laskerian for all $i \geq 0$.

Example 2.5. i) Let $\mathfrak{a}$ be an ideal of R and M an R -module with $\text{Supp}_R M \subseteq V(\mathfrak{a})$. If M is weakly Laskerian, then by Lemma 2.2 ii), it turns out that M is $\mathfrak{a}$ -weakly cofinite. In particular, if either M is finitely generated or Artinian, then M is $\mathfrak{a}$ -weakly cofinite.

ii) Every $\mathfrak{a}$ -cofinite module is $\mathfrak{a}$ -weakly cofinite.

We collect some important known properties of cofinite modules in a lemma. Then we study the corresponding properties for weakly cofinite modules.

Lemma 2.6. i) Suppose M is a finitely generated R -module with $\text{Supp}_R M \subseteq V(\mathfrak{a})$. Then M is $\mathfrak{a}$ -cofinite.

ii) If $0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$ is an exact sequence of R -modules and two of the modules in the exact sequence are $\mathfrak{a}$ -cofinite, then so is the third one.

iii) If M is a $\mathfrak{a}$ -cofinite R -module, then $\text{Ass}_R M$ is finite.

iv) For every $\mathfrak{a}$ -cofinite R -module M , the R -module $M/\mathfrak{a}M$ is finitely generated.

Proof. i) is clear by the definition of cofiniteness.

ii) follows easily from the long Ext sequence induced by the exact sequence

$$0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0,$$

and the fact that if, in an exact sequence, two of the modules in the exact sequence are Noetherian, then the third one is also Noetherian.

iii) Since $\text{Supp}_R M \subseteq V(\mathfrak{a})$, it follows that $\text{Ass}_R M = \text{Ass}_R(0 :_M \mathfrak{a})$. As M is $\mathfrak{a}$ -cofinite, the module $(0 :_M \mathfrak{a}) \cong \text{Ext}_R^0(R/\mathfrak{a}, M)$ is finitely generated. Thus $\text{Ass}_R M$ is finite.

iv) See [13, Corollary 1.2]. $\square$

Remark 2.7. i) Example 2.5 i) shows that the analogue of Lemma 2.6 i) holds for weak cofiniteness. Also, in view of Lemma 2.2 i), it follows that the analogue of Lemma 2.6 ii) holds for weak cofiniteness.

ii) Because the set of associated primes of a weakly Laskerian module is finite, by using the same argument as in the proof of Lemma 2.6 iii), we can deduce that the set of associated primes of a $\mathfrak{a}$ -weakly cofinite module is finite.

iii) Recall that a module M is said to have finite Goldie dimension if M does not contain an infinite direct sum of nonzero submodules, or equivalently, the injective envelope $E(M)$ of M decomposes as a finite direct sum of indecomposable injective submodules. By [13, Proposition 1.3], the Goldie dimension of an $\mathfrak{a}$ -cofinite module is finite. This is not the case for $\mathfrak{a}$ -weakly cofinite modules. To this end, take $M = \bigoplus_{i \in \mathbb{N}} R/\mathfrak{m}$, where $\mathfrak{m}$ is a fixed maximal ideal of R . Since $\text{Supp}_R M = \{\mathfrak{m}\}$ by Lemma 2.2 iii), it turns out that M is weakly Laskerian. Therefore M is $\mathfrak{m}$ -weakly cofinite, although its Goldie dimension is not finite.

Next, we are going to prove the analogue of Lemma 2.6 iv) for weak cofiniteness, but we first need some preliminary results. In view of Lemma 2.2, by using the same proof as that used in [4, Proposition 1], we can deduce the following.

Lemma 2.8. *Let $\mathfrak{a}$ be an ideal of R and N an R -module. For any given integer n , the following are equivalent:*

- i) $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is weakly Laskerian for all $i \leq n$.
- ii) $\text{Ext}_R^i(M, N)$ is weakly Laskerian for all finitely generated R -module M with $\text{Supp}_R M \subseteq V(\mathfrak{a})$, and all $i \leq n$.

The Change of Ring Principle for cofiniteness was proved by Delfino and Marley (see [4, Proposition 2]). In the sequel, we prove the Change of Ring Principle for weak cofiniteness. The proof is an adaption of the proof of [4, Proposition 2].

Theorem 2.9. *Let the ring T be a homomorphic image of R . Let $\mathfrak{a}$ be an ideal of R and M a T -module. Then M is $\mathfrak{a}$ -weakly cofinite as an R -module if and only if M is $\mathfrak{a}T$ -weakly cofinite as a T -module.*

Proof. Assume $T = R/I$ for some ideal I of R and let N be a T -module. Then we have

$$\text{Ass}_T N = \{\mathfrak{p}/I : \mathfrak{p} \in \text{Ass}_R N\}$$

and

$$\text{Supp}_T N = \{\mathfrak{p}/I : \mathfrak{p} \in \text{Supp}_R N\}.$$

Thus N is weakly Laskerian as a T -module if and only if it is weakly Laskerian as an R -module. Also, we deduce that $\text{Supp}_R M \subseteq V(\mathfrak{a})$ if and only if $\text{Supp}_T M \subseteq V(\mathfrak{a}T)$. By [15, Theorem 11.65], there is a Grothendieck spectral sequence

$$E_2^{p,q} = \text{Ext}_T^p(\text{Tor}_q^R(T, R/\mathfrak{a}), M) \xRightarrow[p]{\quad} \text{Ext}_R^{p+q}(R/\mathfrak{a}, M).$$

First assume that M is $\mathfrak{a}T$ -weakly cofinite. Because $\text{Supp}_T(\text{Tor}_q^R(T, R/\mathfrak{a})) \subseteq V(\mathfrak{a}T)$, by Lemma 2.8, it turns out that $E_2^{p,q}$ is a weakly Laskerian T -module for all p and q . For each $r \geq 2$, $E_r^{p,q}$ is a subquotient of $E_2^{p,q}$. Hence, by Lemma 2.2 i) $E_r^{p,q}$ is a weakly Laskerian T -module for all $r \geq 2$ and all $p, q \geq 0$. There is an integer $r \gg 0$ such that $E_\infty^{p,q} = E_r^{p,q}$ for all $p, q \geq 0$. Also, for each $n \in \mathbb{N}_0$ there is a bounded filtration

$$0 = \phi^{n+1}H^n \subseteq \phi^n H^n \subseteq \cdots \subseteq \phi^1 H^n \subseteq \phi^0 H^n = H^n$$

for the module $H^n = \text{Ext}_R^n(R/\mathfrak{a}, M)$ such that $E_\infty^{p,n-p} \cong \phi^p H^n / \phi^{p+1} H^n$ for all $p = 0, 1, \dots, n$. Now, by using Lemma 2.2 i) successively, we deduce that $\text{Ext}_R^n(R/\mathfrak{a}, M)$ is a weakly Laskerian R -module for all $n \geq 0$. That is M is $\mathfrak{a}$ -weakly cofinite.

Conversely, assume that M is $\mathfrak{a}$ -weakly cofinite. By induction on n , we show that $E_2^{n,0} = \text{Ext}_T^n(T/\mathfrak{a}T, M)$ is a weakly Laskerian T -module for all $n \geq 0$. Since the module

$$\text{Hom}_T(T/\mathfrak{a}T, M) \cong \text{Hom}_R(R/\mathfrak{a}, M)$$

is weakly Laskerian, the claim holds for $n = 0$. Now, assume that $n > 0$ and that $E_2^{p,0}$ is weakly Laskerian for all $p < n$. By Lemma 2.8, it turns out that $E_2^{p,q}$ is weakly Laskerian for all $p < n$ and $q \geq 0$. We have $E_r^{n,0} \cong E_\infty^{n,0}$ for all $r \gg 0$. As $H^n = \text{Ext}_R^n(R/\mathfrak{a}, M)$ is weakly Laskerian, it follows that for each $p = 0, 1, \dots, n$, $E_\infty^{p,n-p}$ is weakly Laskerian. From the induction hypothesis, we get that $\text{im } d_{r-1}^{n-r+1, r-2}$ is

weakly Laskerian. Hence $\ker d_{r-1}^{n,0}$ is weakly Laskerian. By continuing this argument, we deduce that $E_2^{n,0} = \ker d_2^{n,0}$ is weakly Laskerian. $\square$

Now, we are ready to present the last result of this section. Its proof is a slight modification of the proof of [13, Corollary 1.2]

Theorem 2.10. *Let $\mathfrak{a}$ be an ideal of R and M a $\mathfrak{a}$ -weakly cofinite R -module. Then $M/\mathfrak{a}M$ is weakly Laskerian.*

Proof. Let $a_1, a_2, \dots, a_n$ be generators of $\mathfrak{a}$. Let $\Theta : R[X_1, X_2, \dots, X_n] \longrightarrow R$ be the natural ring homomorphism defined by $\Theta(f) = f(a_1, a_2, \dots, a_n)$. Then Θ is surjective and $(X_1, X_2, \dots, X_n)R = \mathfrak{a}$. Denote the ideal $(X_1, X_2, \dots, X_n)$ by $\mathfrak{b}$. By Theorem 2.9, $\text{Ext}_{R[X_1, X_2, \dots, X_n]}^i(R[X_1, X_2, \dots, X_n]/\mathfrak{b}, M)$ is weakly Laskerian $R[X_1, X_2, \dots, X_n]$ -module for all $i \geq 0$. Let $K_\bullet(X_1, X_2, \dots, X_n)$ denote the Koszul complex of the ring $R[X_1, X_2, \dots, X_n]$ with respect to $X_1, X_2, \dots, X_n$. Since $X_1, X_2, \dots, X_n$ is a regular sequence in the ring $R[X_1, X_2, \dots, X_n]$, it follows that $K_\bullet(X_1, X_2, \dots, X_n)$ is a free resolution for $R[X_1, \dots, X_n]/\mathfrak{b}$. Thus

$$\text{Ext}_{R[X_1, X_2, \dots, X_n]}^n(R[X_1, X_2, \dots, X_n]/\mathfrak{b}, M) \cong M/\mathfrak{b}M.$$

Therefore $M/\mathfrak{a}M$ is a weakly Laskerian R -module. $\square$

3. ASSOCIATED PRIMES OF LOCAL COHOMOLOGY MODULES

In this section, we prove that the local cohomology modules of a weakly Laskerian module are weakly cofinite in several cases.

Let M be a finitely generated R -module and t a nonnegative integer. In [12, Proposition 2.5], it is shown that if $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -cofinite for all $i \neq t$, then $H_{\mathfrak{a}}^t(M)$ is also $\mathfrak{a}$ -cofinite. In the following we show that the same result holds for weak cofiniteness. Then, as a result we deduce several results concerning finiteness of associated primes of local cohomology modules.

Theorem 3.1. *Let $\mathfrak{a}$ be an ideal of R and M a weakly Laskerian module. Suppose there exists an integer $t \geq 0$ such that $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i \neq t$. Then $H_{\mathfrak{a}}^t(M)$ is also $\mathfrak{a}$ -weakly cofinite.*

Proof. By [15, Theorem 11.38], there is a spectral sequence

$$E_2^{p,q} := \text{Ext}_R^p(R/\mathfrak{a}, H_{\mathfrak{a}}^q(M)) \Longrightarrow_p \text{Ext}_R^{p+q}(R/\mathfrak{a}, M).$$

For each $r \geq 2$, we consider the exact sequence

$$0 \longrightarrow \ker d_r^{p,t} \longrightarrow E_r^{p,t} \xrightarrow{d_r^{p,t}} E_r^{p+r,t-r+1}. (*)$$

It follows from the hypotheses that the R -module $E_r^{p+r,t-r+1}$ is weakly Laskerian. Note that $E_r^{p,q}$ is a subquotient of $E_2^{p,q}$ for all $p, q \in \mathbb{N}_0$.

There is an integer s such that $E_\infty^{p,q} = E_r^{p,q}$ for all $r \geq s$. Also, for each $n \in \mathbb{N}_0$ there is a bounded filtration

$$0 = \phi^{n+1}H^n \subseteq \phi^n H^n \subseteq \cdots \subseteq \phi^1 H^n \subseteq \phi^0 H^n = H^n$$

for the module $H^n = \text{Ext}_R^n(R/\mathfrak{a}, M)$ such that $E_\infty^{p,n-p} \cong \phi^p H^n / \phi^{p+1} H^n$ for all $p = 0, 1, \dots, n$. Thus $E_\infty^{p,q}$ is weakly Laskerian for all p, q . Since

$$E_s^{p,t} = \ker d_{s-1}^{p,t} / \text{im } d_{s-1}^{p-s+1,t+s-2},$$

it follows that $\ker d_{s-1}^{p,t}$ is weakly Laskerian. Hence by using the exact sequence $(*)$ for $r = s-1$, we deduce that $E_{s-1}^{p,t}$ is weakly Laskerian. By continuing this argument repeatedly for integers $s-1, s-2, \dots, 3$ instead of s , we obtain that $E_2^{p,t}$ is weakly Laskerian for all $p \geq 0$. This completes the proof. $\square$

By using the above result, we can deduce the following corollary.

Corollary 3.2. *Let $\mathfrak{a}$ be an ideal of a local ring $(R, \mathfrak{m})$ and M a finitely generated R -module. Assume that $\mathfrak{a}$ contains an M -filter regular sequence $x_1, \dots, x_t$ and that $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i > t$. Then $H_{\mathfrak{a}}^t(M)$ is $\mathfrak{a}$ -weakly cofinite.*

Proof. By [14, Theorem 3.1], $H_{\mathfrak{a}}^i(M)$ is Artinian for all $i < t$. Hence, the claim follows by Lemma 2.2 iii) and Theorem 3.1. $\square$

The following extends the main result of [9].

Corollary 3.3. *Let $\mathfrak{a}$ be a principal ideal of R and M a weakly Laskerian module. Then $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i \geq 0$.*

Proof. Since $H_{\mathfrak{a}}^0(M)$ is a submodule of M , it turns out that $H_{\mathfrak{a}}^0(M)$ is weakly Laskerian, by Lemma 2.2 i). Also, $H_{\mathfrak{a}}^i(M) = 0$ for all $i > 1$. Hence $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i \neq 1$. Therefore, the claim follows by Theorem 3.1. $\square$

Corollary 3.4. *Let $(R, \mathfrak{m})$ be a local ring of dimension d . Let $\mathfrak{a}$ be an ideal of R .*

- a) For any R -module M , the modules $H_{\mathfrak{a}}^d(M)$ and $H_{\mathfrak{a}}^{d-1}(M)$ are weakly Laskerian.*
- b) Assume that M is a weakly Laskerian R -module. The following assertions hold.*
 - i) if $d \leq 3$, then $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i \geq 0$.*

ii) if $d = 4$, then $H_{\mathfrak{a}}^1(M)$ is $\mathfrak{a}$ -weakly cofinite if and only if $H_{\mathfrak{a}}^2(M)$ is $\mathfrak{a}$ -weakly cofinite.

iii) if $d = 4$ and the $\mathfrak{a}$ -ideal transform module of M is weakly Laskerian, then $H_{\mathfrak{a}}^i(M)$ is $\mathfrak{a}$ -weakly cofinite for all $i \geq 0$.

Proof. a) is immediate, because by [11, Corollaries 2.3 and 2.4] $\text{Supp}_R H_{\mathfrak{a}}^d(M)$ and $\text{Supp}_R H_{\mathfrak{a}}^{d-1}(M)$ are finite. Now, we prove b).

i) and ii) are clear by Theorem 3.1 and a).

iii) Let $D_{\mathfrak{a}}(M)$ denote the $\mathfrak{a}$ -ideal transform module of M . There is an exact sequence

$$0 \longrightarrow H_{\mathfrak{a}}^0(M) \longrightarrow M \longrightarrow D_{\mathfrak{a}}(M) \longrightarrow H_{\mathfrak{a}}^1(M) \longrightarrow 0.$$

Hence by Lemma 2.2 i), $H_{\mathfrak{a}}^0(M)$ and $H_{\mathfrak{a}}^1(M)$ are weakly Laskerian. On the other hand, by a) $H_{\mathfrak{a}}^4(M)$ and $H_{\mathfrak{a}}^3(M)$ are both weakly Laskerian. Therefore the conclusion follows, by Theorem 3.1. $\square$

Example 3.5. i) Let $(R, \mathfrak{m})$ be a local ring of dimension 3 and let x, y, z be a system of parameters of R . Let $\mathfrak{a} = (xz, yz)$. Then, by [13, Theorem 2.2], $H_{\mathfrak{a}}^2(R)$ is not $\mathfrak{a}$ -cofinite. But $H_{\mathfrak{a}}^2(R)$ is $\mathfrak{a}$ -weakly cofinite, by Corollary 3.4 i).

ii) Let R and $\mathfrak{a}$ be either as in Example 3.7 or as in Example 3.10 in [12]. Then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^2(R))$ is not finitely generated. Thus $H_{\mathfrak{a}}^2(R)$ is not $\mathfrak{a}$ -cofinite, while in each case $H_{\mathfrak{a}}^i(R)$ is $\mathfrak{a}$ -weakly cofinite for all $i \geq 0$, by Corollary 3.4 i).

iii) Let k be a field, $R = k[x, y][[u, v]]$, $\mathfrak{a} = (u, v)$, and $f = ux + vy$. Then $H_{\mathfrak{a}}^2(R/(f))$ is not $\mathfrak{a}$ -cofinite (see [7, §3]). However, by Corollary 3.4 i), $H_{\mathfrak{a}}^i(R/(f))$ is $\mathfrak{a}(R/(f))$ -weakly cofinite for all $i \geq 0$. In particular, it follows that the set of associated primes of $H_{\mathfrak{a}}^i(R/(f))$ is finite for all $i \geq 0$.

iv) Let $R = \mathbb{Z}[x, y, z, u, v, w]$ and $f = ux + vy + wz$. For the ideal $\mathfrak{a} = (x, y, z)$, Singh [16] has shown that the set of associated primes of $H_{\mathfrak{a}}^3(R/(f))$ is not finite. Thus $H_{\mathfrak{a}}^3(R/(f))$ is not $\mathfrak{a}$ -weakly cofinite, by Remark 2.7 ii).

The following extends [4, Theorem 1] in some sense.

Theorem 3.6. *Let $(R, \mathfrak{m})$ be a local ring and $\mathfrak{a}$ an ideal of R with $\dim R/\mathfrak{a} \leq 1$. Let M be an R -module. Then $H_{\mathfrak{a}}^i(M)$ is weakly Laskerian for all $i \geq 0$.*

Proof. Since $\dim R/\mathfrak{a} \leq 1$, it follows that $V(\mathfrak{a})$ is contained in the set of minimal prime ideals of $\mathfrak{a}$ union with $\{\mathfrak{m}\}$. Thus

$$\text{Supp}_R(H_{\mathfrak{a}}^i(M)) \subseteq V(\mathfrak{a}) \subseteq \text{Ass}_R R/\mathfrak{a} \cup \{\mathfrak{m}\},$$

for all $i \geq 0$. Hence, by Lemma 2.2 iii) $H_{\mathfrak{a}}^i(M)$ is weakly Laskerian. Therefore, the assertion follows. $\square$

The following extends [1, Theorem 1.2].

Theorem 3.7. *Let $(R, \mathfrak{m})$ be a local ring and M a Cohen-Macaulay R -module. Suppose that for any ideal $\mathfrak{a}$ of R , with $t = \text{grade}(\mathfrak{a}, M)$, $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is weakly Laskerian. Then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is a finite set for any ideal $\mathfrak{a}$ of R and all $i \geq 0$.*

Proof. Let $i \geq 0$ be a fixed integer and $\mathfrak{a}$ an ideal of R . We show that the set of associated primes of the local cohomology module $H_{\mathfrak{a}}^i(M)$ is a finite set. If $i \leq \text{grade}(\mathfrak{a}, M)$, then the claim follows by [2, Theorem 2.2]. Hence assume that $i > \text{grade}(\mathfrak{a}, M)$. We may and do assume that $H_{\mathfrak{a}}^i(M) \neq 0$. By [1, Lemma 2.4], there exists an ideal $\mathfrak{b} \supseteq \mathfrak{a}$ of R such that $\text{grade}(\mathfrak{b}, M) = i - 1$ and $H_{\mathfrak{b}}^i(M) \cong H_{\mathfrak{a}}^i(M)$. Now, by the assumption $\text{Hom}_R(R/\mathfrak{b}, H_{\mathfrak{b}}^i(M))$ is weakly Laskerian. Thus $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is a finite set, as required. $\square$

Theorem 3.8. *Let $(R, \mathfrak{m})$ be a local ring and M a Cohen-Macaulay R -module. Let $t \in \mathbb{N}_0$ be a fixed integer such that $H_{\mathfrak{a}}^t(M)$ is weakly Laskerian for any ideal $\mathfrak{a}$ of R . Then $\text{Ass}_R(H_{\mathfrak{a}}^{t+1}(M))$ is a finite set for any ideal $\mathfrak{a}$ of R .*

Proof. Let $\mathfrak{a}$ be an ideal of R . As in the proof of Theorem 3.7, we may assume that $t + 1 \geq \text{grade}(\mathfrak{a}, M)$. Also, we can and do assume that $H_{\mathfrak{a}}^{t+1}(M) \neq 0$. Since M is a Cohen-Macaulay R -module, it follows by [1, Lemma 2.4], that there exists an ideal $\mathfrak{b} \supseteq \mathfrak{a}$ of R such that $\text{grade}(\mathfrak{b}, M) = t$ and $H_{\mathfrak{b}}^{t+1}(M) \cong H_{\mathfrak{a}}^{t+1}(M)$. By the assumption the module $H_{\mathfrak{b}}^t(M)$ is weakly Laskerian. Thus, it follow from [6, Theorem 2.5] that $\text{Ass}_R(H_{\mathfrak{a}}^{t+1}(M))$ is a finite set. $\square$

REFERENCES

- [1] J. Asadollahi and P. Schenzel, *Some results on associated primes of local cohomology modules* Japan. J. Math. (N.S.) **29**(2) (2003), 285-296.
- [2] M.P. Brodmann and F.A. Lashgari, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc., **128**(10) (2000), 2851-2853.
- [3] M.P. Brodmann and R.Y. Sharp: *‘Local cohomology-An algebraic introduction with geometric applications’*, Cambr. Univ. Press, 1998.
- [4] D. Delfino and T. Marley, *Cofinite modules and local cohomology*, J. Pure Appl. Algebra, **121**(1), (1997), 45-52.

- [5] K. Divaani-Aazar and M.A. Esmkhani, *Artinianness of local cohomology modules of ZD-modules*, Comm. Algebra, to appear.
- [6] K. Divaani-Aazar and A. Mafi, *Associated prime of local cohomology modules*, Proc. Amer. Math. Soc., to appear.
- [7] R. Hartshorne, *Affine duality and cofiniteness*, Invent. Math., **9** (1970), 145-164.
- [8] C. Huneke, *Problems on local cohomology*, Free resolutions in commutative algebra and algebraic geometry, Res. Notes Math., **2** (1992), 93-108.
- [9] K.I. Kawasaki, *Cofiniteness of local cohomology modules for principal ideals*, Bull. London Math. Soc., **30**(3) (1998), 241-246.
- [10] K. Khashyaranesh and Sh. Salarian, *On the associated primes of local cohomology modules*, Comm. Algebra, **27**(12) (1999), 6191-6198.
- [11] T. Marley, *The associated primes of local cohomology modules over rings of small dimension*, Manuscripta Math., **104**(4) (2001), 519-525.
- [12] T. Marley and J.C. Vassilev, *Cofiniteness and associated primes of local cohomology modules*, J. Algebra **256**(1) (2002), 180-193.
- [13] L. Melkersson, *Properties of cofinite modules and applications to local cohomology*, Math. Proc. Cambridge Philos. Soc., **125**(3) (1999), 417-423.
- [14] L. Melkersson, *Some applications of a criterion for Artinianness of a module*, J. Pure Appl. Algebra **101**(3) (1995), 291-303.
- [15] J. Rotman, *Introduction to homological algebra*, Academic Press, 1979.
- [16] A.K. Singh, *p-torsion elements in local cohomology modules*, Math. Res. Lett., **7**(2-3) (2000), 165-176.
- [17] A.K. Singh and I. Swanson, *Associated primes of local cohomology modules and of Frobenius powers*, International Mathematics Research Notices **33** (2004) 1703-1733.
- [18] H. Zöschinger, *Minimax-moduln*, J. Algebra **102**(1) (1986), 1-32.

K. DIVAANI-AAZAR, DEPARTMENT OF MATHEMATICS, AZ-ZAHRA UNIVERSITY, VANAK, POST CODE 19834, TEHRAN, IRAN-AND-INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, P.O. BOX 19395-5746, TEHRAN, IRAN.

E-mail address: kdivaani@ipm.ir

A. MAFI, INSTITUTE OF MATHEMATICS, UNIVERSITY FOR TEACHER EDUCATION, 599 TALEGHANI AVENUE, TEHRAN 15614, IRAN.